References – Continence Care Tools

- AGREE Next Steps Consortium. (2009). The AGREE II Instrument [Electronic version].

 Retrieved June 8, 2016 from http://www.agreetrust.org
- Appleyard, J. & Bastone, S. (2015). Redesigning continence services: Rehabilitation and cost-effectiveness. *Journal of Community Nursing*, 29(3), 45-53.
- Association of Rehabilitation Nurses. (2014). *Standards & scope of rehabilitation nursing* practice. (6th ed.). Chicago, IL.
- Ayaş, S., Leblebici, B., Sözay, S., Bayramoğlu, M. & Niron, E.A. (2006) The effect of abdominal massage on bowel function in patients with spinal cord injury. *American Journal of Physical Medicine and Rehabilitation*, 85(12), 951-955. doi:10.1097/01.phm.0000247649.00219.c0
- Barrie, M. (2016). An hour-a-day project aimed at streamlining continence services. *Journal of Community Nursing*, 30(2), 45-53. Retrieved from http://www.jcn.co.uk/files/downloads/articles/50945-jcn-april-may-13-streamlining-continence-services1.pdf
- Bayliss, V., Cherry, M., Lock, R. & Salter, L. (2000). Pathways for continence care: development of the pathways. *British Journal of Nursing*, *9*(17), 1165-1172.
- Bayliss, V. & Salter, L. (2004). Pathways for evidence-based continence care. *Nursing Standard*, 19(9), 45-51.
- Booth, J. Kumlien, S., Zang, Y., Gustafsson, B. & Tolson, D. (2009). Rehabilitation nurses practices in relationship to urinary incontinence following stroke: a cross-cultural

- comparison. *Journal of Clinical Nursing*, *18*, 1049-1058. doi: 10.1111/j.1365-2702.2008.02688.x
- Borrie, M.J., Campell, K., Arcese, Z.A., Bray, J., Hart, P. and Labate, T. (2001) Urinary retention in patients in a geriatric rehabilitation unit: Prevalence, risk factors, and validity of bladder scan evaluation. Rehabilitation Nursing, 26, 187-191.
- Boyer, D.R. & Stelzer, N. (2009). Implementation of an evidence-based bladder scanner protocol. *Journal of Nursing Care Quality*, 24(10), 10-16.
- Brooks, W. (2004). The use of practice guidelines for urinary incontinence following stroke.

 *British Journal of Nursing, 13(20), 1176-1179.
- Callan, L.L. & Willson, M. (2016). Fecal incontinence: Pathology, assessment and management.

 In D.B. Doughty & K.N. Moore (Eds.), *Core Curriculum: Continence Management*.(pp. 297-327). Philadelphia, PA: Wolters Kluwer.
- Cameron, A.P., Rodriguez, G.M., Gursky, A., He, C., Clemens, J.Q. & Stofeel, J.T. (2015). The severity of bowel dysfunction in patients with neurogenic bladder. *The Journal of Urology*, 194, 1336-1341. doi:10.1016.j.juro.2015.04.100
- Cave, C. (2016). Evidence based continence care: An integrative review. *Rehabilitation Nursing*, 0, 1-14. doi:10.1002/rnj.291
- Callan, L.L. & Willson, M. (2016). Fecal incontinence: Pathology, assessment and management.

 In D.B. Doughty & K.N. Moore (Eds.), *Core Curriculum: Continence Management*.(pp. 297-327). Philadelphia, PA: Wolters Kluwer.
- Chipps, T. (2011). Using behavioral methods to manage faecal incontinence. *Britsh Journal of Nursing*, 20(8), 1172-1178.

- Coffey, A., McCarthy, G., McCormack, B., Wright, J. & Slater, P. (2007). Incontinence:

 Assessment, diagnosis and management in two rehabilitation units for older people.

 Worldviews on Evidence-Based Nursing. 4(4), 179-186.
- Coggrave, M.J., Burrows, D., & Durand, M.A. (2006). Progressive protocol in the bowel management of spinal cord injuries. *British Journal of Nursing*, *15*(20), 1108-1113.
- Coggrave, M.J. & Norton, C. (2009). The need for manual evacuation and oral laxatives in the management of neurogenic bowel dysfunction after spinal cord injury: a randomized controlled trial of a stepwise protocol. *Spinal Cord*, 48, 504-510. doi:10.1038/sc.2009.166
- Coggrave, M., Norton, C., & Cody, J.D. (2014). Management of faecal incontinence and constipation in adults with central neurological diseases (Review). The Cochrane Collaboration. John Wiley & Sons, Ltd. doi:10.1002/14651858.CD002115.pub5
- Cooper, G. & Watt, E. (2003). An exploration of acute care nurses' approach to assessment and management of people with urinary incontinence. *Journal of Wound Ostomy and Continence Nursing*, 30(6), 305-313. doi: 10.1016/mjw.2003.157
- Cournan, M. (2012). Bladder management in female stroke survivors: translating research to practice. *Rehabilitation Nursing*, *37*(5), 220-230. doi: 10.1002/rnj.054
- Davies, E.C., Green, C.F., Mottram, D.R., & Pirmohamed, M. (2008). The use of opioids and laxatives, and incidence of constipation, in patients requiring neck-of-femur (NOF) surgery: A pilot study. *Journal of Clinical Pharmacy and Therapeutics*, 33, 561-566.

- Denby, N. (2006). The role of diet and lifestyle changes in the management of constipation. *Journal of Community Nursing*, 20(9), 20-9.
- Doughty, D.B. & Moore, K.N. (2016). Overview of urinary incontinence and voiding dysfunction. In D.B. Doughty & K.N. Moore (Eds.), *Core Curriculum: Continence Management*.(pp. 297-327). Philadelphia, PA: Wolters Kluwer.
- Dowling-Castronovo, A. & Specht, J.K. (2009). Assessment of transient urinary incontinence in older adults. *American Journal of Nursing*, 109(2), 62-71.
- Dumoulin, C., Korner-Bitensky, N., Tannenbaum, C. (2005). Urinary continence after stroke:

 Does rehabilitation make a difference? A systemic review of the evidence of behavior therapy. *Topics in Stroke Rehabilitation*, 12(3), 66-76.
- Emmanuel, A. I (2010). Managing neurogenic bowel dysfunction. *Clinical Rehabilitation*, 24, 483-488. doi:10.1177/0269215509353253
- Ersoz, M., et al. (2013). An evaluation of bladder emptying methods and the effect of demographic and clinical factors on spontaneous voiding frequency in stroke patients.

 Neurological Science, 34, 729-734. doi:10.1007/s10072-012-1123-9
- Fowler, C.J., Griffiths, D. & de Groat, W.C. (2008). The neural control of micturition. *National Review of Neuroscience*, 9(6), 453-466. doi:10.1038/nrn2401
- Grandstaff, M. & Lyons, D. (2012). Impact of a continence training program on patient safety and quality. *Rehabilitation Nursing*, *37*(4), 180-184.
- Gross, J.C. (2003). Urinary incontinence after stroke: Evaluation and behavioral treatment. *Topics in Geriatric Rehabilitation*, 19(1), 60-83.

- Harris, J., Roussel, L., Dearman, C., & Thomas, P. (2016). Project planning and management: A guide for nurses and interprofessional teams (2nd ed). Sudbury, MA:Jones and Bartlett Publishing.
- Herr-Wilbert, I., Imhof, L., Hund-Georgiadis, M., & Wilbert, D. (2010). Assessment-guided therapy of urinary incontinence after stroke. *Rehabilitation Nursing*, 35(6), 248-253.
- Hsu, C.C. & Sandofrd, B.A. (2007). The Delphi technique: Making sense of consensus.

 *Practical Assessment, Research & Evaluation, 12(10), 1-7.
- Jacelon, C. (Ed.). (2011). The specialty practice of rehabilitation nursing: A core curriculum. (6th ed.). Glenview, IL: Association of Rehabilitation Nurses.
- King, L. & Pilcher, M. (2008). A multidisciplinary approach to improving urinary continence.

 *Nursing Standard, 23(8), 42-46.
- Leary, S.M., Liu, C., Cheesman, A.L., Ritter, A., Thompson, S. & Greenwood, R. (2006).

 Incontinence after brain injury: Prevalence and multidisciplinary management on a neurological rehabilitation unit. *Clinical Rehabilitation*, 20, 1094-1099.
- Lehman, C.A. & Owen, S.V. (2001). Bladder scanner accuracy during everyday use on an acute rehabilitation unit. *Spinal Cord Injury Nursing*, *18*(2), 87-92.
- Linari, L.R., Schofield, L.C. & Horrom, K.A. (2011). Implementing a bowel program.

 Orthopedic Nursing, 30(5), 317-321. doi: 10.1097/NOR.0b013e31822c5c10
- Lim, S.F. & Childs, C. (2013). A systematic review of the effectiveness of bowel management strategies for constipation in adults with stroke. *International Journal of Nursing Studies*, 50, 1004-1010. doi:http://dx.doi.org/10.1016.j.ijnurstu.2012.12.002

- Matthews, M. (2010). Causes and rehabilitation of urinary incontinence after stroke: a literature review. *British Journal of Neuroscience Nursing*, 6(1), 37-41.
- McClurg, D., Hagen, S., Hawkins, S. & Lowe-Strong, A. (2011). Abdominal massage for the alleviation of constipation symptoms in people with multiple sclerosis: A randomized controlled feasibility study. *Multiple Sclerosis Journal*, *17*(2), 223-233. doi:10.1177/13524585110384899
- Mehdi, Z., Birns, J. & Bhalla, A. (2013). Post-stroke urinary incontinence. *International Journal of Clinical Practice*, 67(11), 1128-1137. doi:10.111/jjcp.12183
- Milne, J. (2008). Behavioral therapies for overactive bladder: Making sense of the evidence. *Journal of Wound, Ostomy and Continence Nursing*, 35(1), 93-101.
- Nazarko, L. (2010). Assessing and treating bladder problems after stroke. *Nursing and Residential Care*, 12(4), 182-185.
- Nazarko, L. (2015). Person-centered care of women with urinary incontinence. *Nurse Prescribing*, *13*(6), 288-293.
- Newman, D.K. & Wein, A.J. (2009). *Managing and treating urinary incontinence* (2nd ed).

 Baltimore, MD: Health Professions Press.
- Norton, C. & Chelvanayagam, S. (2010). Bowel problems and coping strategies in people with multiple sclerosis. *British Journal of Nursing*, 19(4), 220-226.
- O'Connell, B., Ostaszkiewicz, J., & Hawkins, M. (2011). Innovations in aged care: A suite of evidence-based continence assessment tools for residential aged care. *Australian Journal of Ageing*, 30(1), 27-32. doi:10.1111/j.1741-6612.2010.004499.x

- Ostaszkiewicz, J. (2006). A clinical nursing leadership model for enhancing continence care for older adults in a subacute inpatient care setting. *Journal of Wound, Ostomy and Continence Nursing*, 33(6), 624-629.
- Ostaszkiewicz, J., O'Connell, B., & Millar, L. (2008). Incontinence: Managed or mismanaged in hospital settings? *International Journal of Nursing Practice*, *14*, 495-502. doi:10.1111/j.1440-172X.2008.00725.x
- Ostaszkiewicz, J., Ski, C. & Hornby, L. (2005). Does successful treatment of constipation or faecal impaction resolve lower urinary tract symptoms? A structured review of the literature. *Australian and New Zealand Continence Journal*, 11(3), 70-80.
- Pellatt, G.C. (2008). Neurogenic Continence. Part 1: Pathophysiology and quality of life. *British Journal of Nursing*, *17*(13), 836-841.
- Pellatt, G.C. & Geddis, T. (2008). Neurogenic continence. Part 2: Neurogenic bladder management. *British Journal of Nursing*, *17*(14), 904-913.
- Pryor, J. & Jannings, W. (2005). Promoting self-efficacy in bowel care following spinal cord injury: A pilot project. *Journal of the Australasian Rehabilitation Nurses' Association*, 8(4), 21-25.
- Rao, S.S.C. & Go, J.T. (2010). Update on the management of constipation in the elderly: New treatment options. *Clinical Interventions in Aging*, *5*, 163-171.
- Richmond, J.P. & Wright, M.E. (2006). Development of a constipation risk assessment scale. *Journal of Orthopedic Nursing*, 19, 186-197. doi:10.1016/j.joon.2006.08.001

- Rigby, D. (2002). The value of continence training: does it change clinical practice? *British Journal of Nursing*, 12(8), 484-492.
- Roe, B., Flanagan, L. & Maden, M. (2015). Systematic review of systematic reviews for the management of urinary incontinence and promotion of continence using conservative behavioral approaches in older people in care homes. *Journal of Advanced Nursing*, 71(7), 1464-1482. doi:10.1111/jan.12613
- Roe, B., Ostaszkiewicz, J., Milne, J. & Wallace, S. (2006). Systematic reviews of bladder training and voiding programmes in adults: A synopsis of findings from data analysis and outcomes using metastudy techniques. *Journal of Advanced Nursing*, *57*(1), 15-31. doi:10.1111/j.1365-2648.2006.04097.x
- Rigby, D. (2003). The value of continence training: does it change clinical practice? *British Journal of Nursing*, 12(8), 484-492.
- Schnelle, J.F., et al. (2010). A controlled trial of an intervention to improve urinary and fecal incontinence and constipation. *Journal of American Geriatric Society*, 58, 1504-1511.
- Selius, B.A. and Subedi, R. (2008) Urinary retention in adults: Diagnosis and initial management. *American Family Physician*, 77(5), 643-650. Retrieved on December 18, 2016 from http://www.aafp.org/afp/2008/0301/p643.html
- Shalmani, H.M., et al. (2010). Direct and indirect costs of functional constipation: A population-based study. *International Journal of Colorectal Diseases*, 26, 515-522. doi:10.1007/s00384-010-1077-4
- Smith, G.D. & Watson, R. (2008). Gastrointestinal Nursing. Oxford, UK. Blackwell Publishing

- Solomons, J. & Woodward, S. (2013). Digital removal of faeces in the bowel management of patients with spinal cord injury: A review. *British Journal of Neuroscience Nursing*, 9(5), 216-222.
- Sparks, et al. (2004). The clinical benefits of the bladder scanner: A research synthesis. *Journal of Nursing Care Quality*, 19(3), 188-192.
- Steele, S.E., Han, C.J. & Heitkemper. (2016). Motiliy disorders. In D.B. Doughty & K.N. Moore (Eds.), *Core Curriculum: Continence Management*. (pp. 297-327). Philadelphia, PA: Wolters Kluwer.
- Testa, A. (2015). Understanding urinary incontinence in adults. *Urologic Nursing*, 35(2), 82-86. doi:10.7257/1053-816X.2015.35.2.82
- Thomas, et al., (2014). Identifying continence options after stroke (ICONS): A cluster randomized controlled feasibility trial. *Trials Journal*, *15*(509), 1-15. Retrieved on June 1, 2016 from http://trialsjournal.biomedcentral.com/articles/10.1186/1745-6215-12-131
- Tod, A.M., Stringer, E., Levery, C., Dean, J. & Brown, J. (2007). Rectal irrigation in the management of functional bowel disorders: A review. *British Journal of Nursing*, *16*(14), 858-864.
- Vaughn, S. (2009). Efficacy of urinary guidelines in the management of post-stroke incontinence. *International Journal of Urological Nursing*, 3(1), 4-12.
- Vaughn, S., Mauk, K., Jacelon, C., Larsen, P., Rye, J. ... Dufresne, D. (2015). The competency model for professional rehabilitation nursing. *Rehabilitation Nursing Journal*, 41, 33-44. doi: 10.1002/rnj.225
- Ward-Smith, P. (2009). The cost of urinary incontinence. *Urologic Nursing*, 29(3), 188-194.

- Washington University in St Louis. (2012). The program sustainability assessment tool [Webbased instrument]. Center for Public Health System Science: George Warren Brown School of Social Work. Retrieved from https://sustaintool.org/
- Weiss, B.D. (2005). Selecting medications for the treatment of urinary incontinence. *American Family Physician*, 7192), 315-322. Retrieved on November 21, 2016 from http://www.aafp.org/afp/2005/0115/p315.pdf
- Wilde, M.H., Bliss, D.A., Booth, J., Cheater, F.M. & Tannenbaum (2014). Self-management of urinary and fecal incontinence. *American Journal of Nursing* 114(1), 38-45.
- Wilson, L. (2005). Understanding bowel problems in older people: part 2. *Nursing Older People*, 17(9), 24-29.
- Williams, K., Assassa, R., Smith, N., Rippin, C., Shaw, C. & Mayne, C. (2002). Good practice in continence care development for nurse-led service. *British Journal of Nursing*, 11(8), 548-559.
- Woodward, S. (2013). Managing urinary incontinence in people with neurological disorders: Part 2: Interventions. *British Journal of Neuroscience Nursing*, *9*(2), 63-70.
- Woodward, S. (2014). Managing urinary incontinence in people with neurological disorders.

 *Nursing and Residential Care, 16(6), 319-324.

 doi:http://dx.doi.org/10.12968/nrec.2014.16.6.319

- Wright, J., McCormack, B., Coffey, A. & McCarthy. (2006). Developing a tool to assess personcentered continence care. *Nursing Older People*, *18*(6), 23-28.
- Wright, J., McCormack, B., Coffey, A. & McCarthy. (2007). Evaluating the context within which continence care is provided in rehabilitation units for older people. *International Journal of Older People Nursing*, 2(1), 9-19.
- Yap, T.L., Cromwell, D.C. & Emberton, M. (2006). A systematic review of the reliability of frequency-volume charts in urological research and its implications for the optimum chart duration. *BJU International*, *99*, 9-16, doi:10.1111/j.1464-410X.2006.06499.x
- Yu, N.S., Pomeranz, J.L., Moorhouse, M.D., Shaw, L.R., & Deutsch, P.M. (2008). Identifying a new area of damages: Assessing time loss association with bowel management. *Journal of Life Care Planning*, 7(1), 3-11.